



Confidence Interval for the Difference of 2 Population Means

Ha Khanh Nguyen (hknguyen)

X_1, X_2, \dots, X_{n_1}
from population 1
mean μ_1 , std σ_1

Y_1, Y_2, \dots, Y_{n_2}
from population 2
mean μ_2 , std σ_2

When one of these 2 conditions are satisfied:

- n_1 and n_2 are large (≥ 30)
- Population 1 and population 2 are approximately normal

Then, $(\bar{X} - \bar{Y})$ is approximately normal with mean $\mu_1 - \mu_2$ and standard deviation $\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$.

A $(1 - \alpha)100\%$ confidence interval for $\mu_1 - \mu_2$ is

$$(\bar{X} - \bar{Y}) \pm z_{\alpha/2} \cdot \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$$

If σ_1 and σ_2 are unknown, use:

$$(\bar{X} - \bar{Y}) \pm t_{\alpha/2} \cdot \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

where the number of degrees of freedom of the t -dist = smaller of $n_1 - 1$ and $n_2 - 1$.

Notes: If n_1 and n_2 are large, $t_{\alpha/2}$ can be approximated by $z_{\alpha/2}$.

Example 1:

Dr. Statman claims that his new revolutionary study method “Study While You Sleep” (SWYS) is more effective than the traditional study methods. In an experiment, 250 students enrolled in the same section of STAT 400 at UIUC were divided into 2 groups. 100 students volunteered to study using SWYS method, and the other 150 students did whatever the students usually do. At the end of the semester, the average of the total number of points (out of 500) were compared for the 2 groups.

Note: This is not a good experiment design!

	SWYS	Traditional
Average total points	450	410
Standard deviation	20	45

Construct a 95% confidence interval for the difference in the average total points for SWYS and the traditional study methods.

Step 0: Check the assumptions

Step 1: Compute the $(1 - \alpha)100\%$ confidence interval